

PRINCIPAL-VALUE RESUMMATION FOR DILEPTON PRODUCTION

HARRY CONTOPANAGOS

*High Energy Physics Division, Argonne National Laboratory,
Argonne, IL 60439-4815, USA*

and

LYNDON ALVERO, GEORGE STERMAN

*Institute for Theoretical Physics, SUNY Stony Brook,
Stony Brook, NY 11794-3840, USA*

ABSTRACT

We present a new method of resummation of QCD threshold distributions to the hard scattering function of the dilepton production cross section in hadron-hadron collisions. Our formulation by-passes the infrared singularities of the QCD running coupling through a principal value prescription, and does not require an explicit infrared cut-off. The resulting large corrections exponentiate, and we discuss asymptotic properties of the exponent. We present a closed analytical formula for the resummed hard part in momentum space, and give predictions, at fixed-target energies, for the resulting cross section which is a bounded function of its kinematic variables in the entire partonic phase space.

1. Introduction

It is well-known that perturbative QCD (pQCD) corrections to certain hadron-hadron scattering cross sections can be numerically important even at quite high energies. One source of large corrections is soft-gluon radiation near partonic threshold. The best-studied example of this mechanism is the dilepton-production cross section, which is not only of great phenomenological importance but is also a true laboratory for testing various ideas on pQCD. As we shall see, at fixed-target energies, the resummed series can differ significantly from even two-loop expansions. Indeed, the perturbative series evaluated order-by-order does not appear to converge, and in fact, due to the presence of infrared (IR) renormalons, is at best only asymptotic. This suggests that

predictions of the normalization of such cross sections on the basis of low-order calculations combined with parton distributions determined from deeply inelastic scattering are unreliable.

Resummation formulas typically organize the large perturbative corrections in an exponential containing integrals of the QCD running coupling $\alpha_s(\mu^2)$ over various momentum scales. The resulting expressions, however, suffer from divergences¹ associated with the singularity in the perturbative coupling at Λ_{QCD} . One way to deal with this problem is to introduce an explicit infrared cutoff. The dependence on such a cutoff is higher-twist¹. The resulting dependence on the cutoff, however, is difficult to analyze, and is not negligible in general.

Here we summarize recent work^{2,3,4,5} that advances a new resummation method of large threshold corrections, which we have called principal-value resummation. In this approach, the integral over the momentum scales of the running coupling in the resummed exponent is defined as a principal value integral, thus by-passing the IR singularities of the coupling. It can therefore be taken as a mathematically unambiguous definition of the resummed perturbative series. This then allows us to define and explore the structure of its higher-twist corrections, since they are defined relative to a well-defined perturbative expression. Furthermore, the new resummed exponent exhibits the behavior anticipated in pQCD: at each point in partonic phase space it is best-approximated by a particular polynomial in α_s . It also becomes a purely non-perturbative, higher-twist expression in the non-perturbative region of the partonic phase space. The resulting hard part is calculable numerically as well as analytically (including inversion of the Mellin transform). We will summarize all of the above properties within the context of the Drell-Yan process.

2. Principal-Value Resummation

We will focus on the reaction

$$h_1(p_1) + h_2(p_2) \rightarrow l\bar{l}(Q) + X, \quad (1)$$

where in the following we use the notation $s = (p_1 + p_2)^2$, and $z = \tau/x_a x_b$, with x_a, x_b the momentum fractions of the scattered partons. We will concentrate on the diagonal-flavor quark-antiquark subprocess which is singular in the soft-gluon limit $z \rightarrow 1$ at fixed order of pQCD.

The corresponding cross section may be written as

$$\frac{d\sigma}{dQ^2} = \sigma_0^V(Q^2) W^V(\tau, Q^2), \quad (2)$$

where σ_0^V is the point-like tree-level cross section for $q\bar{q} \rightarrow V^* \rightarrow l\bar{l}$, containing all electroweak parameters and physical dimensions and

$$W^V(\tau, Q^2) = \sum_f C_f^V \int_0^1 \frac{dx_a}{x_a} \frac{dx_b}{x_b} F_{f/h_1}(x_a, Q^2) F_{\bar{f}/h_2}(x_b, Q^2) \omega_{f\bar{f}}(\tau/x_a x_b, \alpha), \quad (3)$$

where $\alpha \equiv \alpha_s(Q^2)/\pi$ and C_f^V , in units of $|e|$, is the coupling constant of parton f to vector boson V . Specifically, $C_f^\gamma = e_f^2$ and $C_f^Z = 1 + (1 - 4|e_f| \sin^2 \theta_W)^2$. The sum in

eq. (3) runs over all relevant quark and antiquark flavors and the parton distributions F_{f/h_1} , $F_{\bar{f}/h_2}$ must be in DIS scheme form.

For the Mellin transform of the hard part, $\omega_{f\bar{f}}(z, \alpha)$, we obtain the principal-value resummation formula in moment space³

$$\tilde{\omega}_{f\bar{f}}(n, \alpha) = A(\alpha)e^{E(n, \alpha)}, \quad (4)$$

where the function $A(\alpha)$ exponentiates large “constants”⁶, i.e., terms that depend only on α and not on the parton momenta, and the resummation exponent $E(n, \alpha)$ depends on the soft gluon momenta through the correspondence $n \leftrightarrow 1/(1-z)$. It is this quantity that we will define in principal-value resummation.

We may write¹

$$\Gamma_{f\bar{f}}(1-\zeta, Q^2) \equiv \int_0^\zeta \frac{dy}{1-y} g_1(\alpha[(1-\zeta)(1-y)Q^2]) + g_2(\alpha[(1-\zeta)Q^2]), \quad (5)$$

and define the principal-value exponent as³

$$E(n, Q^2) = - \int_P d\zeta \frac{\zeta^{n-1} - 1}{1-\zeta} \Gamma_{f\bar{f}}(1-\zeta, Q^2), \quad (6)$$

where $P \equiv \frac{1}{2} \times C_+ \cup \frac{1}{2} \times C_-$, with $C_+(C_-)$ any contour in the ζ plane connecting 0 and 1 above (below) the real axis and with $\alpha(\lambda Q^2)$ truncated to only those terms that create the large threshold corrections,

$$\alpha(\lambda Q^2) \equiv \alpha(\lambda) = \frac{\alpha}{1 + \alpha b_2 \ln \lambda} - \frac{\alpha^2 b_3 \ln(1 + \alpha b_2 \ln \lambda)}{b_2 (1 + \alpha b_2 \ln \lambda)^2}, \quad (7)$$

where $b_2 = \frac{1}{12}(33 - 2n_f)$ and $b_3 = \frac{1}{48}(306 - 38n_f)$, n_f being the number of active flavors. The functions g_i , $i = 1, 2$ in eq. (5) are simple expansions of the form $g_i(\alpha(\lambda)) = \sum_{j=1}^\infty \alpha^j(\lambda) g_i^{(j)}$ and the corresponding coefficients are factorization scheme- and process-dependent and can be fitted to finite-order results. What makes the procedure finite and useful is that we *only need* three of these coefficients to account for *all* large threshold corrections to all orders. These, in the DIS scheme, are

$$g_1^{(1)} = 2C_F, \quad g_2^{(1)} = -\frac{3}{2}C_F, \quad g_1^{(2)} = C_F \left[C_A \left(\frac{67}{18} - \frac{1}{6}\pi^2 \right) - \frac{5}{9}n_f \right]. \quad (8)$$

It can be shown⁴ that the *purely* perturbative regime is defined, in moment space, by the relation

$$1 < \ln n < \frac{1}{2\alpha b_2} \leftrightarrow 1 \ll n < \frac{Q}{\Lambda}, \quad (9)$$

where, in the second relation, we have used for simplicity a 1-loop relation between α and $\Lambda \equiv \Lambda_{QCD}$. In this perturbative range, a small- α asymptotic approximation leads to the perturbative formula for the exponent

$$E(n, \alpha) \simeq E(n, \alpha, \mathcal{N}) = \sum_{i=1}^2 \sum_{\rho=1}^{N_i+1} \alpha^\rho \sum_{j=0}^{\rho+1} s_{j,\rho}(i) \ln^j n + \sum_{i=3}^7 \sum_{\rho=1}^{N_i+1} \alpha^\rho \sum_{j=0}^{\rho} s_{j,\rho}(i) \ln^j n, \quad (10)$$

where the $\{s_{j,\rho}(i)\}$ are calculable numerical coefficients and $\mathcal{N} = \mathcal{N}(Q) = \{N_i\}$, the set of optimum numbers of perturbative terms, is determined by optimization of the perturbative asymptotic approximation, eqs. (6), (10). At typical fixed-target dilepton mass scales, say $Q = 10$ GeV, we find^{3,4} $\mathcal{N} = \{2, 6, 6, 2, 6, 1, 6\}$.

It can similarly be shown⁴ that the *purely* non-perturbative regime is defined through the relation

$$\ln n > \frac{1}{\alpha b_2} \leftrightarrow n > \frac{Q^2}{\Lambda^2}. \quad (11)$$

Exact analytical results for the exponent in this region have been obtained⁴ through a large- α asymptotic approximation, but suffice it to say that for extremely large values of n ,

$$E(n, \alpha) \simeq -\frac{g_1^{(1)}}{b_2} \ln 2 \ln n - \frac{g_2^{(1)}}{b_2} \ln(\alpha b_2 \ln n) - \frac{g_1^{(1)} b_3}{4b_2^3} \ln^2(\alpha b_2 \ln n). \quad (12)$$

This shows that the exponent tends to $-\infty$ at the edge of phase space $z \simeq 1$ (*i.e.*, $n \rightarrow \infty$). As a result, the corresponding resummed cross section is finite, because the hard part is exponentially damped at that threshold. This results in the principal-value cross section being a bounded function of τ throughout its kinematic range, in contrast to the finite-order cross section that diverges as a power of the logarithm as $\tau \rightarrow 1$.

To obtain the resummed hard part in momentum space, we perform an inverse Mellin transform on eq. (4) analytically². The final result for the resummed hadronic function, eq. (3), is

$$W^V(\tau, Q^2) = \sum_f C_f^V A(\alpha) \int_\tau^1 dz [1 + \mathcal{H}(z, \alpha)] \frac{d}{dz} \left(\frac{\mathcal{F}_{f\bar{f}}(\tau/z)}{z} \right), \quad (13)$$

where the parton flux is defined as

$$\mathcal{F}_{f\bar{f}}(\tau/z) = \int_0^1 \delta(1 - \tau/(zx_a x_b)) \frac{dx_a}{x_a} \frac{dx_b}{x_b} F_{f/h_1}(x_a, Q^2) F_{\bar{f}/h_2}(x_b, Q^2), \quad (14)$$

and the hard function $\mathcal{H}$ is given by^{2,4}

$$\mathcal{H}(z, \alpha) \equiv \int_0^z dz' \frac{e^{E(\frac{1}{1-z'}, \alpha)}}{\pi(1-z')} \Gamma \left(1 + P_1 \left(\frac{1}{1-z'}, \alpha \right) \right) \sin \left(\pi P_1 \left(\frac{1}{1-z'}, \alpha \right) \right), \quad (15)$$

where Γ is the Euler Gamma function and $P_1(x, \alpha) \equiv \frac{\partial}{\partial \ln x} E(x, \alpha)$. This hard function may be used for predicting both the mass-distribution for the process and its derivative with respect to rapidity or x_F ⁵.

3. Comparison with experiment and conclusions

Because of space limitation, we will treat only a single example: the E537 fixed target experiment at Fermilab⁷, which observed Drell-Yan pairs produced by antiprotons on a tungsten target at $\sqrt{s} = 15.3$ GeV ($V = \gamma$ in eqs. (2) and (3)). In fig. 1, we plot the data as well as the one-, two-loop and resummed predictions for the mass

distribution $Q^3 d\sigma/dQ$ integrated over $x_F > 0$. We used a parton distribution⁸ based on a fit to the *purely* DIS data of CDHS. Notice that agreement between the resummation prediction and the data is acceptable, and better than that of the finite-order prediction. Parton distributions with larger sea-quark content tend to give larger predictions in the resummed cross section.

We conclude that principal-value resummation is an appropriate way to organize consistently large threshold corrections in pQCD to all orders in perturbation theory, and that experimental results can be successfully described in this framework. This also suggests that whatever higher-twist part of the theory is not contained within pQCD may be suppressed, even in the domain of phase space probed by fixed-target experiments at momenta like those cited in fig. 1. The inclusion of the information contained in resummation formulas may also be important for the determination of parton distributions by the combination of DIS and hadron-hadron data.




Fig. 1. E537 experiment. Short-dash: 1-loop; Long-dash: 2-loop; Solid: Resummed.

Acknowledgements

Supported by the U.S. Department of Energy, Division of High Energy Physics, contract W-31-109-ENG-38, and in part by the National Science Foundation under grant PHY 9309888 and the Texas National Research Laboratory.

References

1. G. Sterman, *Nucl. Phys.* **B281** (1987) 310;
D. Appell, P. Mackenzie and G. Sterman, *Nucl. Phys.* **B309** (1988) 259.
2. H. Contopanagos and G. Sterman, *Nucl. Phys.* **B400** (1993) 211.

This contains a full list of references.

3. H. Contopanagos and G. Sterman, *Nucl. Phys.* **B419** (1994) 77.
4. L. Alvero and H. Contopanagos, ANL-HEP-PR-94-25/ITP-SB-94-33 (July 1994).
5. L. Alvero and H. Contopanagos, *in preparation*.
6. L. Magnea and G. Sterman, *Phys. Rev.* **D42** (1990) 4222.
7. E. Anassontzis et al., *Phys. Rev. Lett.* **54** (1985) 2572.
8. B. Betev et al, *Z. Phys.* **C28** (1985) 15; J.S. Conway et al, *Phys. Rev.* **D39** (1989) 92; K. Freudenreich, *Int. Jour. of Mod. Phys.* **A5** (1990) 3643.